
Radiofrequency Ablation of Kidney Tumors

This information is reviewed by a physician with expertise in the area presented and is further reviewed by committees from the American College of Radiology (ACR) and the Radiological Society of North America (RSNA), comprising physicians with expertise in several radiologic areas.

What is Radiofrequency Ablation of Kidney Tumors?

Radiofrequency ablation, sometimes referred to as RFA, is a minimally invasive treatment for cancer. It is an image-guided technique that heats and destroys cancer cells.

In radiofrequency ablation, imaging techniques such as ultrasound, computed tomography (CT) or magnetic resonance imaging (MRI) are used to help guide a needle electrode into a cancerous tumor. High-frequency electrical currents are then passed through the electrode, creating heat that destroys the abnormal cells.

What are some common uses of the procedure?

Radiofrequency ablation is used to treat renal cell carcinoma (kidney tumors).

Radiofrequency ablation is a viable and effective treatment option if you:

- have one kidney.
- have other medical conditions which might prevent surgery.
- are older and might have difficulty with surgery or postsurgical recovery.
- have tumors of less than four centimeters in size.
- have a familial predisposition (family history) to multiple kidney tumors.

RFA may also be used pre-operatively to decrease blood loss during surgery.

How should I prepare?

You may be instructed not to eat or drink anything after midnight before your procedure. Your doctor will tell you which medications you may take in the morning.

You should report to your doctor all medications that you are taking, including herbal supplements, and if you have any allergies, especially to local anesthetic medications, general anesthesia, or to contrast materials (also known as “dye” or “x-ray dye”). Your physician may advise you to stop taking aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs) or a blood thinner for a specified period of time before your procedure.

Prior to your procedure, your blood may be tested to determine how well your liver and kidneys are functioning and whether your blood clots normally.

Women should always inform their physician or x-ray technologist if there is any possibility that they are pregnant. Many imaging tests are not performed during pregnancy so as not to expose the fetus to radiation. If an x-ray is necessary, precautions will be taken to minimize radiation exposure to the baby.

You should plan to have a relative or friend drive you home after your procedure.

You will be given a gown to wear during the procedure.

What does the equipment look like?

In this procedure, computed tomography (CT), needle electrodes, an electrical generator and grounding pads may be used.

There are two types of needle electrodes: simple straight needles and a straight, hollow needle that contains several retractable electrodes that extend when needed.

The radiofrequency generator produces electrical currents in the range of radiofrequency waves. It is connected by insulated wires to the needle electrodes and to grounding pads that are placed on the patient's back or thigh.



The CT scanner is typically a large, box like machine with a hole, or short tunnel, in the center. You will lie on a narrow examination table that slides into and out of this tunnel. Rotating around you, the x-ray tube and electronic x-ray detectors are located opposite each other in a ring, called a gantry. The computer workstation that processes the imaging information is located in a separate room, where the technologist operates the scanner and monitors your examination.

Other equipment used during the procedure includes an intravenous line (IV) and equipment that monitors your heart beat and blood pressure.

How does the procedure work?

Radiofrequency ablation works by passing electrical currents in the range of radiofrequency waves between the needle electrode and the grounding pads placed on the patient's skin. These currents create heat around the electrode, which when directed into the tumor, heats and destroys the cancer cells. Because healthy liver tissue is better able to withstand heat, radiofrequency ablation is able to destroy a tumor and only a small rim of normal tissue around the edges of the tumor. At the same time, heat from radiofrequency energy closes small blood vessels and lessens the risk of bleeding. The dead tumor cells are gradually replaced by scar tissue that shrinks over time.

Ultrasound or computed tomography imaging may be used to help the physician guide the needle electrode into the tumor.

How is the procedure performed?

Image-guided, minimally invasive procedures such as radiofrequency ablation are most often performed by a specially trained interventional radiologist in an interventional radiology suite or occasionally in the operating room.

Radiofrequency ablation is often done on an outpatient basis.

You will be positioned on the examining table.

You will be connected to monitors that track your heart rate, blood pressure and pulse during the procedure.

A nurse or technologist will insert an intravenous (IV) line into a vein in your hand or arm so that sedation medication can be given intravenously.

The area where the electrodes are to be inserted will be sterilized and covered with a surgical drape.

Your physician will numb the area with a local anesthetic.

A very small nick is made in the skin at the site.

If multiple electrodes are needed, then multiple nicks may be made.

Radiofrequency ablation is performed using one of the following methods:

- Surgically.
- Percutaneous, in which needle electrodes are inserted through the skin and into the site of the tumor.

Using imaging guidance, your physician will insert the needle electrode through the skin and advance it to the site of the tumor.

Once the needle electrode is in place, radiofrequency energy is applied. For a large tumor, it may be necessary to do multiple ablations by repositioning the needle electrode into different parts of the tumor to ensure no tumor tissue is left behind.

At the end of the procedure, the needle electrode will be removed and pressure will be applied to stop any bleeding and the opening in the skin is covered with a dressing. No sutures are needed.

Each radiofrequency ablation takes about 10 to 30 minutes, with additional time required if multiple ablations are performed. The entire procedure is usually completed within one to three hours.

What will I experience during and after the procedure?

Devices to monitor your heart rate and blood pressure will be attached to your body.

You will feel a slight pin prick when the needle is inserted into your vein for the intravenous line (IV) and when the local anesthetic is injected.

The intravenous (IV) sedative will make you feel relaxed and sleepy. You may or may not remain awake, depending on how deeply you are sedated.

Pain immediately following radiofrequency ablation can be controlled by pain medication given through your IV or by injection. Afterward any mild discomfort you experience can be controlled by oral pain medications. Patients may feel nauseous, but this can also be relieved by medication.

You will remain in the recovery room until you are completely awake and ready to return home.

Your intravenous line will be removed.

You should be able to resume your usual activities within a few days.

Only about two percent of patients still have pain a week following radiofrequency ablation.

Who interprets the results and how do I get them?

Computed tomography (CT) or magnetic resonance imaging (MRI) of the kidneys is performed within a few hours to a week following radiofrequency ablation. A radiologist will interpret these CT or MRI scans to detect any complications and to ensure that all of the tumor tissue has been destroyed.

What are the benefits vs. risks?

Benefits

- RFA is a relatively quick procedure and recovery is rapid so that chemotherapy may be resumed almost immediately.
- Radiofrequency ablation is less expensive than other treatment options.
- No surgical incision is needed—only a small nick in the skin that does not have to be stitched closed.
- Unlike surgical treatment, this procedure allows the kidney to be preserved.
- It does not have an effect on blood pressure.
- It may or may not have an effect on renal function depending on how large an area needs to be treated.

Risks

- Severe pain after RFA is uncommon, but may last a few days and require a narcotic to provide relief.
- Any procedure where the skin is penetrated carries a risk of infection. The chance of infection requiring antibiotic treatment appears to be less than one in 1,000.
- There is always a slight chance of cancer from excessive exposure to radiation. However, the benefit of this treatment outweighs the risk.
- Women should always inform their physician and x-ray or CT technologist if there is any possibility that they are pregnant.

What are the limitations of Radiofrequency Ablation of Kidney Tumors?

Research indicates that RFA is very effective for small tumors. Because it has only recently begun to be used for kidney tumors, research is limited. Long term effectiveness at five and 10 years are available in only a limited number of patients.

Disclaimer:

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images are shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright © 2009 Radiological Society of North America, Inc.
Send comments via email to: webmast2@rsna.org