



## Ultrasound - Prostate

### What is Ultrasound Imaging of the Prostate?

Ultrasound imaging, also called ultrasound scanning or sonography, involves exposing part of the body to high-frequency sound waves to produce pictures of the inside of the body. Ultrasound examinations do not use ionizing radiation (as used in x-rays). Because ultrasound images are captured in real-time, they can show the structure and movement of the body's internal organs, as well as blood flowing through blood vessels.

Ultrasound imaging is a noninvasive medical test that helps physicians diagnose and treat medical conditions.

Prostate ultrasound, also called transrectal ultrasound, provides images of a man's prostate gland and surrounding tissue. The exam typically requires insertion of an ultrasound probe into the rectum of the patient. The probe sends and receives sound waves through the wall of the rectum into the prostate gland which is situated right in front of the rectum.

### What are some common uses of the procedure?

A transrectal ultrasound of the prostate gland is performed to:

- detect disorders within the prostate.
- determine whether the prostate is enlarged, with measurements acquired as needed for any treatment planning.
- detect an abnormal growth within the prostate.
- help diagnose the cause of a man's infertility.

A transrectal ultrasound of the prostate gland is typically used to help diagnose symptoms such as:

- a nodule felt by a physician during a routine physical exam or prostate cancer screening exam.
- an elevated blood test result.
- difficulty urinating.

Because ultrasound provides real-time images, it also can be used to guide procedures such as needle biopsies, in which a needle is used to sample cells (tissue) from an abnormal area in the prostate gland for later laboratory testing.

## How should I prepare?

You should wear comfortable, loose-fitting clothing for your ultrasound exam. You may need to remove all clothing and jewelry in the area to be examined.

You may be asked to wear a gown during the procedure.

You may be instructed to avoid taking blood thinners, such as aspirin, for seven to 10 days prior to the procedure if a biopsy is planned. An enema may be taken two to four hours before the ultrasound to clean out the bowel.

## What does the equipment look like?

Ultrasound scanners consist of a console containing a computer and electronics, a video display screen and a transducer that is used to do the scanning. The transducer is a small hand-held device that resembles a microphone, attached to the scanner by a cord. The transducer sends out inaudible high frequency sound waves into the body and then listens for the returning echoes from the tissues in the body. The principles are similar to sonar used by boats and submarines.

The ultrasound image is immediately visible on a video display screen that looks like a computer or television monitor. The image is created based on the amplitude (strength), frequency and time it takes for the sound signal to return from the area of the patient being examined to the transducer and the type of body structure the sound travels through.

For ultrasound procedures requiring insertion of the transducer, such as transvaginal or transrectal exams, the device is covered and lubricated.



## How does the procedure work?

Ultrasound imaging is based on the same principles involved in the sonar used by bats, ships, fishermen and the weather service. When a sound wave strikes an object, it bounces back, or echoes. By measuring these echo waves, it is possible to determine how far away the object is and its size, shape and consistency (whether the object is solid, filled with fluid, or both).

In medicine, ultrasound is used to detect changes in appearance of organs, tissues, and vessels or detect abnormal masses, such as tumors.

In an ultrasound examination, a transducer both sends the sound waves and receives/records the echoing waves. When the transducer is pressed against the skin, it directs small pulses of inaudible, high-frequency sound waves into the body. As the sound waves bounce off of internal organs, fluids and tissues, the sensitive microphone in the transducer records tiny changes in the sound's pitch and direction. These signature waves are instantly measured and displayed by a computer, which in turn creates a real-time picture on the monitor. One or more frames of the moving pictures are typically captured as still images. Small loops of the moving “real time” images may also be saved.



The same principles apply to ultrasound procedures such as transrectal and transvaginal which require insertion of a special transducer into the body.

## How is the procedure performed?

In men, the prostate gland is located directly in front of the rectum, so the ultrasound exam is performed transrectally.

For a transrectal ultrasound, you will be asked to lie on your side with your knees bent. A disposable protective cover is placed over the transducer, it is lubricated, inserted through the anus and placed into the rectum.

The images are obtained from different angles to get the best view of the prostate gland.

If a suspicious lesion is identified with ultrasound or with a rectal examination, an ultrasound-guided biopsy can be performed. This procedure involves advancing a needle into the prostate gland while the radiologist watches the needle placement with ultrasound. A small amount of tissue is taken for microscopic examination. Shown is an example of a transrectal transducer (probe).



When the examination is complete, the patient may be asked to dress and wait while the ultrasound images are reviewed.

This ultrasound examination is usually completed in less than 20 minutes.

## What will I experience during and after the procedure?

Ultrasound exams in which the transducer is inserted into an opening of the body may produce minimal discomfort.

If no biopsy is required, transrectal ultrasound of the prostate is similar or may have less discomfort than a rectal exam performed by your doctor.

If a biopsy is performed, additional discomfort, due to the needle insertion, is usually minimal because the rectal wall is relatively insensitive to the pain in the region of the prostate. A biopsy will add time to the procedure.

Rarely, a small amount of blood may be present in the sperm or urine following the procedure.

After an ultrasound examination, you should be able to resume your normal activities immediately.

## Who interprets the results and how do I get them?

A radiologist, a physician specifically trained to supervise and interpret radiology examinations, will analyze the images and send a signed report to your primary care physician or the physician who referred you for the exam, who will share the results with you. In some cases the radiologist may

discuss results with you at the conclusion of your examination.

Follow-up examinations are often necessary, and your doctor will explain the exact reason why another exam is requested. Sometimes a follow-up exam is done because a suspicious or questionable finding needs clarification with additional views or a special imaging technique. A follow-up examination may be necessary so that any change in a known abnormality can be detected over time. Follow-up examinations are sometimes the best way to see if treatment is working or if an abnormality is stable over time.

## What are the benefits vs. risks?

### Benefits

- Ultrasound is widely available, easy-to-use and less expensive than other imaging methods.
- Ultrasound imaging uses no ionizing radiation.
- Ultrasound scanning gives a clear picture of soft tissues that do not show up well on x-ray images.
- Ultrasound causes no health problems and may be repeated as often as is necessary if medically indicated.
- Ultrasound provides real-time imaging, making it a good tool for guiding minimally invasive procedures such as needle biopsies and needle aspiration.

### Risks

- For standard diagnostic ultrasound there are no known harmful effects on humans.

## What are the limitations of Prostate Ultrasound Imaging?

Men who have had the tail end of their bowel (rectum) removed during prior surgery are not good candidates for ultrasound of the prostate gland because this type of ultrasound typically requires placing a probe into the rectum.

## Additional Information and Resources

### RadiologyInfo

Prostate Cancer  
([www.RadiologyInfo.org/en/info.cfm?pg=pros\\_cancer](http://www.RadiologyInfo.org/en/info.cfm?pg=pros_cancer))

### RTAnswers.org

Radiation Therapy for Prostate Cancer  
([www.rtanswers.com/treatmentinformation/cancertypes/prostate/index.aspx](http://www.rtanswers.com/treatmentinformation/cancertypes/prostate/index.aspx))

## Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

## Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2011 Radiological Society of North America, Inc.