

Vascular Access Procedures

This information is reviewed by a physician with expertise in the area presented and is further reviewed by committees from the American College of Radiology (ACR) and the Radiological Society of North America (RSNA), comprising physicians with expertise in several radiologic areas.

What are Vascular Access Procedures?

A vascular access procedure involves the insertion of a flexible thin plastic tube, or catheter, into a blood vessel to provide a painless way of drawing blood or delivering drugs and nutrients into a patient's bloodstream over a period of weeks, months or even years.

A simple intravenous (IV) line is effective for short-term use, but is not suitable for long-term use. When an IV line is necessary for a longer period of time and/or a more secure venous access is necessary, a special catheter, called a central access catheter, or a similar device is placed inside a major blood vessel either temporarily (days) or long-term (weeks to years) so that it can be easily and repeatedly accessed over a desired period of time.

In a vascular access procedure, a special catheter is inserted inside a major vein (generally in one of the large veins in the neck or in the arm) extending into the large central vein near the heart.

What are some common uses of the procedure?

Vascular access procedures are performed when patients need:

- intravenous antibiotic treatment.
- chemotherapy, or anti-cancer drugs.
- long-term intravenous (IV) feeding for nutritional support.
- repeated drawing of blood samples.
- hemodialysis, a process used to treat patients whose kidneys are not working properly. It involves a special machine and tubing that removes blood from the body, cleanses it of waste and extra fluid and then returns it back to the body.

Access catheters may also be used for:

- blood transfusions.
- patients who have difficulty receiving a simple IV line.

How should I prepare for the procedure?

Prior to your procedure, your blood may be tested to determine how well your liver and kidneys are functioning and whether your blood clots normally.

You should report to your doctor all medications that you are taking, including herbal supplements, and if you have any allergies, especially to local anesthetic medications, general anesthesia, or to contrast materials (also known as "dye" or "x-ray dye"). Your physician may advise you to stop taking aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs) or a blood thinner for a specified period of time days before your procedure.

Women should always inform their physician or x-ray technologist if there is any possibility that they are pregnant. Many imaging tests are not performed during pregnancy so as not to expose the fetus to radiation. If an x-ray is necessary, precautions will be taken to minimize radiation exposure to the baby.

You will receive specific instructions on how to prepare, including any changes that need to be made to your regular medication schedule.

You may be instructed not eat or drink anything for several hours before your procedure.

You should plan to have a relative or friend drive you home after your procedure.

You may be asked to remove some or all of your clothes and to wear a gown during the exam. You may also be asked to remove jewelry, eye glasses and any metal objects or clothing that might interfere with the x-ray images.

What does the equipment look like?

In this procedure, x-ray and ultrasound equipment, a vascular access catheter, a venous access needle and a guide wire are used.

The equipment typically used for this examination consists of a radiographic table, an x-ray tube and a television-like monitor that is located in the examining room or in a nearby room. When used for viewing images in real time, the image intensifier (which converts x-rays into a video image) is suspended over a table on which the patient lies. When used for taking still pictures, the image is captured either electronically or on film.

The x-ray equipment allows the operator to watch the wire and catheter on a live display so they can be inserted safely and positioned to optimize catheter function.

Ultrasound scanners consist of a console containing a computer and electronics, a video display screen and a transducer that is used to scan the body.

Ultrasound is used to assess the vein that is being used for catheter placement and to provide guidance during the venous puncture. It is important to use ultrasound guidance during the venous puncture to reduce possible complications such as bleeding.

In contrast to the catheter used in a standard intravenous (IV) line, a vascular access catheter is more durable and does not easily become blocked or infected. These catheters are designed in a way that they extend into the largest central vein near the heart.

A catheter is a long, thin plastic tube, about as thick as a strand of spaghetti. Following are the major types of vascular access catheters:

- The peripherally inserted central catheter (PICC) is a long catheter that extends from an arm vein into the largest vein (superior vena cava) near the heart and typically provides central IV access for several weeks, but may remain in place for several months. These catheters are called “midline catheters” when they are placed in a way that the tip of the catheter remains in a relatively large vein, but doesn’t extend into the largest central vein. They may have one or two lumens and some may be able to be used for CT contrast injections (manufactured for forceful contrast injections).
- Non-tunneled central catheters are larger caliber than PICC, and they are designed to be placed via a relatively large, more central vein such as the jugular vein in the neck or the femoral vein in the groin.

- The tunneled catheter has a cuff that stimulates tissue growth that will help hold it in place in the body. Examples of the tunneled catheter include HICKMAN® catheters, BROVIAC® catheters and GROSHONG® catheters. There are several different types of dialysis catheters. The tunneled catheter is the best choice when access to the vein is needed for long period of time. It is secure and easy to access. They are more secure and usually work more efficiently than PICCs because of their design (cuff on the catheter stimulating tissue growth) and larger size.

The port catheter, or subcutaneous implantable port, is a permanent device that consists of a catheter attached to a small reservoir, both of which are placed under the skin similar to tunnel catheters.

Note: HICKMAN®, BROVIAC® and/or GROSHONG® are registered trademarks of C. R. Bard, Inc. and its related company, BCR, Inc.

A small, hollow needle and long thin wire, called a guide wire, help the physician position the catheter.

Other equipment used during the procedure includes an intravenous line (IV) and equipment that monitors your heart beat and blood pressure.

How is the procedure performed?

A midline catheter and some peripherally inserted central catheter (PICC) lines may be inserted at your bedside without image guidance. These are inserted through a vein near the elbow and threaded through a large vein in the upper arm.

Other vascular access procedures are most often performed by a specially trained interventional radiologist in an interventional radiology suite or occasionally in the operating room.

These procedures are often done on an outpatient basis.

You will be positioned on your back. A nurse or technologist may insert an intravenous (IV) line into a vein in your hand or arm so that sedative medication can be given intravenously. PICC placement usually does not require sedative medications.

The area of your body where the catheter is to be inserted will be shaved (typically the upper chest for tunneled catheters and in the arm for PICCs), sterilized and covered with a surgical drape.

Your physician will numb the area with a local anesthetic. A very small nick in the skin is made at the site.

PICC:

To place a PICC line, the physician will identify the vein using ultrasound or x-ray guidance and insert a small needle into the arm vein and advance a small guide wire into the large central vein, called the superior vena cava, under x-ray (fluoroscopy). The catheter is then advanced over the guide wire and moved into position. The guide wire is then removed. If this is done without x-ray guidance, a chest x-ray is needed to confirm the catheter position.

NON-TUNNELED CENTRAL CATHETERS:

These catheters are placed via a relatively larger vein such as the jugular vein in the neck or femoral vein in the groin.

TUNNELED CATHETERS:

For a tunneled catheter, the physician will make one small nick in the skin commonly in the lower neck. Using ultrasound guidance, access is gained into the vein, usually the jugular vein at the base of the neck, and a small guide wire is advanced into the large central vein, called the superior vena cava, under x-ray guidance (fluoroscopy). A subcutaneous tunnel is then created. Using x-ray guidance, the catheter is placed through the tunnel into the superior vena cava. The cuff, which is typically made of Dacron®, is located on the tunneled part of the catheter. Finally, the physician will place stitches at the end of the tunnel to help keep the catheter firmly in place.

PORT-CATHETERS:

Implanting a subcutaneous port, generally requires two incisions (except in the arm where a single incision may suffice). The port reservoir is placed under the skin following a small skin incision and creating a small subcutaneous pocket. The incision for the port is usually about two inches long. The rest of the procedure is similar to the tunneled central catheter placement. A small, elevated area remains on your body at the site of the reservoir. The port, which passes from an access site in a vein of your arm, shoulder or neck, ends in a large central vein in the chest. The reservoir has a silicone covering that can be punctured with a special needle.

Incisions are held together by stitches, surgical glue and/or a special tape.

An x-ray may be performed after the procedure to ensure the catheter is positioned correctly.

Your intravenous line will be removed.

The implanted vascular access catheter is then ready for use.

What will I experience during the procedure?

Devices to monitor your heart rate and blood pressure will be attached to your body.

You will feel a slight pin prick when the needle is inserted into your vein for the intravenous line (IV) and when the local anesthetic is injected.

The intravenous (IV) sedative will make you feel relaxed and sleepy. You may or may not remain awake, depending on how deeply you are sedated.

You may feel some pressure or brief discomfort when the needle is placed into the vein and when the tunnel is created. You will have to lay flat for about 30 to 45 minutes during catheter placement.

If you are not staying overnight at the hospital, you should rest at home for the remainder of the day following the procedure. You may resume your usual activities the next day, but should avoid lifting heavy objects.

After having a tunneled catheter or subcutaneous port placed, you may experience bruising, swelling and tenderness in the chest, neck or shoulder, but these symptoms clear up in a few days. Pain medicine may help during this time.

You will receive instructions on how to care for your incision(s) and your particular vascular access device. For the first week, it is especially important to keep the catheter site clean and dry. Some, but not all, physicians will recommend sponge bathing around the catheter site, then cleaning the area with peroxide, applying an anesthetic ointment that contains an antibiotic and bandaging the area.

You may be allowed to shower after one week, using a piece of plastic wrap over the site where the catheter was inserted. You should not allow the incision to be held under water by swimming or soaking in a tub.

You may be advised to flush your catheter with a heparin solution to help keep blood clots from forming and blocking the catheter.

You should call the physician or nurse if you have any questions about your vascular access device or if:

- the device malfunctions.
- there is bleeding at the insertion site or signs of infection.
- you develop a fever.

- you notice redness, increased swelling, tenderness, warmth or fluid drainage at the catheter insertion site.

Who interprets the results and how do I get them?

Your interventional radiologist will use x-ray imaging during the placement procedure or a chest x-ray taken immediately after the procedure to confirm that your catheter is correctly positioned. Your physician will also check how well your vascular access device is functioning by using a needle and/or syringe and injecting fluid through the catheter.

What are the benefits vs. risks?

Benefits

- A central catheter permits infusion of solutions containing medication or nutritional substances without causing the complications that may occur with an IV, such as local tissue damage when a toxic drug leaks out of the vein.
- In many conditions, having this type of tube inserted provides a simple and painless means of drawing blood, or delivering drugs, nutrients or both.
- Vascular access devices spare the patient the discomfort and stress of repeated needle sticks.
- The vascular access device is an extremely useful solution for patients who—for any reason—require repeated entry into the venous circulation over a long period. A number of different designs are available that are suitable for different circumstances.
- Placement of a vascular access device is a great solution for those requiring prolonged treatment such as chemotherapy. They will not need to have an IV line placed for each treatment and their arm veins will not become badly scarred.
- A PICC is very helpful when medicines or fluids that are irritating to the wall of the vein are needed. A wide range of products may be given by this route, including antibiotics and blood products, and the catheter also may be used for IV feeding and frequent blood sampling.
- A vascular access device may be used immediately after placement. Some types will continue functioning well for a year or longer. The devices are easily removed when no longer needed.

- A catheter sometimes is the only way of getting access to the circulatory system for hemodialysis in patients with serious kidney disease.

Risks

Two types of risk are associated with vascular access devices: those occurring during or shortly after placement and delayed risks that occur simply because the device is in your body.

Following are some of the risks associated with placement of a vascular access device:

- Any procedure that involves placement of a catheter inside a blood vessel carries certain risks. These risks include damage to the blood vessel, bruising or bleeding at the puncture site and infection.
- An infection may develop at an incision site shortly after catheter placement. The risk is less if you carefully follow instructions about caring for the incisions as they heal.
- Bleeding (hemorrhaging). This risk can be minimized through a blood test in advance to be sure that your blood clots normally. If it does not, the procedure may be postponed or you may receive medication or blood products to improve blood clotting.
- Very rarely a patient may develop a condition called a pneumothorax, a collection of air in the chest that may cause one of the lungs to collapse. This may occur during placement of a catheter or port using a vein in the chest or neck, but not when an arm vein is used. The risk is lessened when catheter placement is guided by ultrasound or fluoroscopy. This is why these catheters should be placed by an interventional radiologist using appropriate imaging guidance during the procedure.
- The normal heart rhythm may be disturbed while the catheter is inserted, but this is usually only temporary. The problem is easily recognized during the procedure and eliminated by adjusting the catheter position.
- Rarely, the catheter will enter an artery rather than a vein. If this happens, the catheter will have to be removed. Most often the artery then heals by itself, but occasionally it has to be surgically repaired.

Delayed Risks

- Two types of delayed infection may develop: skin infection at the catheter or port insertion site or bloodstream infection. Infections are least common after placing a port. The risk of delayed

infection can be minimized if you and anyone else who will be handling the device wash hands before flushing it or cleaning the insertion site. The site should be carefully inspected each time the dressing is changed. The risk of infection is higher for individuals who have low white blood cell counts.

- A hole or break in the catheter may lead to leakage of fluid. Breaks may be avoided by not always clamping the catheter in the same spot and by never using too much force when flushing it. Two important first aid measures: 1) clamp the catheter between the damaged part and the skin insertion site; 2) tape a sterile gauze pad to the skin to cover the break. Catheters rarely fracture inside the body. A chest x-ray will show a fracture and allow removal of the broken fragment without surgery.
- The catheter may become accidentally dislodged. If the catheter is not secured to the skin appropriately, it may come out. If this happens, you should apply pressure to the incision site using a sterile dressing and call your physician immediately.
- Air in the catheter is an emergency that may cause chest pain or shortness of breath. You should clamp the catheter right away, lie on your left side and call 9-1-1. This problem can be avoided by always clamping the catheter before and after inserting a syringe, and by making sure that the catheter cap is screwed on tightly.
- Any type of vascular access catheter may become obstructed by clotted blood or fibrin sheath. You can minimize the risk by carefully following instructions about flushing the catheter. Once a catheter occludes, or becomes closed off, it sometimes can be cleared by injecting medication but at other times must be removed or exchanged for a new catheter. Occasionally the catheter can be stripped by a simple interventional procedure.
- If the vein in which the catheter lies becomes occluded (closed off), the arm, shoulder, neck or head may develop swelling. If this occurs, call your physician immediately. The clot may be treated by a blood-thinning medication, but occasionally the catheter will have to be removed.

The reservoir septum of most types of implanted ports has a useful lifetime of about 1,000 punctures and so is not suitable for patients who require IV access on a daily basis.

Some patients have very poor veins that are not well suited for catheter placement. This usually happens when these access veins have been used for a long period of time (years for TPN, etc.). It may be very difficult to find a suitable vein to place a catheter in these patients, and may require unusual venous entry sites (e.g., through the back or through the liver).

Disclaimer:

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images are shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright © 2009 Radiological Society of North America, Inc. Send comments via email to: webmast2@rsna.org

What are the limitations of Vascular Access Procedures?

Although some types of central venous catheter may remain in place for months or even years, most catheters require replacement after certain time frame because of poor function.