

Abdominal and Pelvic MRI

Magnetic resonance imaging (MRI) of the abdomen and pelvis uses a powerful magnetic field, radio waves and a computer to produce detailed pictures of the inside of your body. It may be used to help diagnose or monitor treatment for a variety of conditions within the abdomen and pelvis. If you're pregnant, MRI may be used to safely monitor you or your baby.

Tell your doctor about any health problems, recent surgeries or allergies and whether there's a possibility you are pregnant. The magnetic field is not harmful, but it may cause some medical devices to malfunction. Most orthopedic implants pose no risk, but you should always tell the technologist if you have any medical devices or metal in your body.

Guidelines about eating and drinking before your exam vary between facilities. Unless you are told otherwise, take your regular medications as usual. Leave jewelry at home and wear loose, comfortable clothing. You may be asked to wear a gown. If you have claustrophobia or anxiety, you may want to ask your doctor to prescribe a mild sedative before the exam.



What is abdominal and pelvic MRI?

Magnetic resonance imaging (MRI) is a noninvasive test doctors use to diagnose medical conditions.

MRI uses a powerful magnetic field, radiofrequency pulses, and a computer to produce detailed pictures of internal body structures. MRI does not use radiation (x-rays).

Detailed MR images allow doctors to examine the body and detect disease.

What are some common uses of the procedure?

MR imaging of the abdomen and pelvis is performed to evaluate:

- organs of the abdomen, such as the liver (<http://www.radiologyinfo.org>) , biliary tract (<http://www.radiologyinfo.org>) , kidneys (<http://www.radiologyinfo.org>) , spleen (<http://www.radiologyinfo.org>) , bowel (<http://www.radiologyinfo.org>) , pancreas (<http://www.radiologyinfo.org>) , and adrenal glands (<http://www.radiologyinfo.org>) .
- organs of the pelvis, such as the bladder (<http://www.radiologyinfo.org>) and the reproductive organs such as the uterus (<http://www.radiologyinfo.org>) and ovaries (<http://www.radiologyinfo.org>) in females and the prostate gland (<http://www.radiologyinfo.org>) in males.
- blood vessels (including MR Angiography (<https://www.radiologyinfo.org/en/info/angiomr>)).
- lymph nodes.

Doctors use MRI to help diagnose or monitor treatment for conditions such as:

- tumors of the abdomen or pelvis.
- diseases of the liver, such as cirrhosis (<https://www.radiologyinfo.org/en/info/cirrhosisliver>) , and abnormalities of the bile ducts and pancreas.

- inflammatory bowel disease such as Crohn's disease (<https://www.radiologyinfo.org/en/info/crohns-disease>) and ulcerative colitis (<http://www.radiologyinfo.org>) .
- abnormal blood vessels and inflamed vessels (vasculitis).
- a fetus in the womb of a pregnant woman.

How should I prepare for the procedure?

You will need to change into a hospital gown. This is to prevent artifacts appearing on the final images and to comply with safety regulations related to the strong magnetic field.

Guidelines about eating and drinking before an MRI vary between specific exams and facilities. Take food and medications as usual unless your doctor tells you otherwise.

Some MRI exams use an injection of contrast material. The doctor may ask if you have asthma or allergies to contrast material, drugs, food, or the environment. MRI exams commonly use a contrast material called gadolinium (<http://www.radiologyinfo.org>) . Doctors can use gadolinium in patients who are allergic to iodine contrast. A patient is much less likely to be allergic to gadolinium than to iodine contrast. However, even if the patient has a known allergy to gadolinium, it may be possible to use it after appropriate pre-medication. *For more information on allergic reactions to gadolinium contrast, please consult the ACR Manual on Contrast Media* (<https://www.acr.org/Clinical-Resources/Contrast-Manual>) .

Tell the technologist or radiologist if you have any serious health problems or recent surgeries. Some conditions, such as severe kidney disease, may mean that you cannot safely receive gadolinium. You may need a blood test to confirm your kidneys are functioning normally.

Women should always tell their doctor and technologist if they are pregnant. MRI has been used since the 1980s with no reports of any ill effects on pregnant women or their unborn babies. However, the baby will be in a strong magnetic field. Therefore, pregnant women should not have an MRI in the first trimester unless the benefit of the exam clearly outweighs any potential risks. Pregnant women should not receive gadolinium contrast unless absolutely necessary. *See the MRI Safety During Pregnancy* (<https://www.radiologyinfo.org/en/info/safety-mri-pregnancy>) page for more information about pregnancy and MRI.

If you have claustrophobia (<http://www.radiologyinfo.org>) (fear of enclosed spaces) or anxiety, ask your doctor to prescribe a mild sedative prior to the date of your exam.

Leave all jewelry and other accessories at home or remove them prior to the MRI scan. Metal and electronic items are not allowed in the exam room. They can interfere with the magnetic field of the MRI unit, cause burns, or become harmful projectiles. These items include:

- jewelry, watches, credit cards, and hearing aids, all of which can be damaged
- pins, hairpins, metal zippers, and similar metallic items, which can distort MRI images
- removable dental work
- pens, pocketknives, and eyeglasses
- body piercings
- mobile phones, electronic watches, and tracking devices.

In most cases, an MRI exam is safe for patients with metal implants, except for a few types. People with the following implants may not be scanned and should not enter the MRI scanning area without first being evaluated for safety:

- some cochlear (ear) implants
- some types of clips used for brain aneurysms (<http://www.radiologyinfo.org>)

- some types of metal coils placed within blood vessels
- some older cardiac defibrillators (<http://www.radiologyinfo.org>) and pacemakers (<http://www.radiologyinfo.org>)
- vagal nerve stimulators

Tell the technologist if you have medical or electronic devices in your body. These devices may interfere with the exam or pose a risk. Many implanted devices will have a pamphlet explaining the MRI risks for that device. If you have the pamphlet, bring it to the attention of the scheduler before the exam. MRI cannot be performed without confirmation and documentation of the type of implant and MRI compatibility. You should also bring any pamphlet to your exam in case the radiologist or technologist has any questions.

If there is any question, an x-ray can detect and identify any metal objects. Metal objects used in orthopedic surgery generally pose no risk during MRI. However, a recently placed artificial joint may require the use of a different imaging exam.

Tell the technologist or radiologist about any shrapnel, bullets, or other metal that may be in your body. Foreign bodies near and especially lodged in the eyes are very important because they may move or heat up during the scan and cause blindness. Dyes used in tattoos may contain iron and could heat up during an MRI scan. This is rare. The magnetic field will usually not affect tooth fillings, braces, eyeshadows, and other cosmetics. However, these items may distort images of the facial area or brain. Tell the radiologist (<http://www.radiologyinfo.org>) about them.

Infants and young children usually require sedation or anesthesia to complete an MRI exam without moving. Whether your child requires sedation depends on their age, intellectual development and the type of exam. Moderate and conscious sedation can be provided at many facilities. A doctor or nurse specializing in sedation or anesthesia for children should be available during the exam for your child's safety. You will be given special instructions for how to prepare your child.

Some hospitals employ certified child life specialists to provide children and families with emotional support in medical settings. These specialists have backgrounds in child development, psychology and counseling. They can prepare children for medical imaging procedures. This can help decrease the child's stress and anxiety and even reduce or eliminate the need for sedation.

Ask your doctor if child life specialists are available at your imaging facility.

Many facilities offer child-friendly imaging suites decorated with murals and lighting that can help entertain and calm pediatric patients. Silent MRI techniques and distraction devices like DVD goggles and music headphones may reduce or eliminate the need for sedation during MRI.

New and improved MRI approaches produce high-quality images and reduce the time children spend in the scanner. This may also eliminate the need for sedation. *For more information, see the Pediatric Sedation (<https://www.radiologyinfo.org/en/info/safety-pediatric-sedation>) page.*

What does the equipment look like?

The traditional MRI unit is a large cylinder-shaped tube surrounded by a circular magnet. You will lie on a table that slides into a tunnel towards the center of the magnet.

Some MRI units, called short-bore systems (<http://www.radiologyinfo.org>), are designed so that the magnet does not completely surround you. Some newer MRI machines have a larger diameter bore, which can be more comfortable for larger patients or those with claustrophobia. "Open" MRI units are open on the sides. They are especially helpful for examining larger patients or those with claustrophobia. Open MRI units can provide high quality images for many types of exams. Open MRI may not be used for certain exams. For more information, consult your radiologist.

How does the procedure work?

Unlike x-ray and computed tomography (CT) exams, MRI does not use radiation. Instead, radio waves re-align hydrogen atoms that naturally exist within the body. This does not cause any chemical changes in the tissues. As the hydrogen atoms return to their usual alignment, they emit different amounts of energy depending on the type of tissue they are in. The scanner captures this energy and creates a picture using this information.

In most MRI units, the magnetic field is produced by passing an electric current through wire coils. Other coils are inside the machine and, in some cases, are placed around the part of the body being imaged. These coils send and receive radio waves, producing signals that are detected by the machine. The electric current does not come into contact with the patient.

A computer processes the signals and creates a series of images, each of which shows a thin slice of the body. The radiologist can study these images from different angles.

MRI is often able to tell the difference between diseased tissue and normal tissue better than x-ray, CT, and ultrasound.

How is the procedure performed?

MRI examinations may be performed on outpatients or inpatients.

The technologist will position you on the moveable exam table. They may use straps and bolsters to help you stay still and maintain your position.

The technologist may place devices that contain coils capable of sending and receiving radio waves around or next to the area of the body under examination.

MRI exams generally include multiple runs (sequences), some of which may last several minutes. Each run will create a different set of noises.

If your exam uses a contrast material, a doctor, nurse, or technologist (<http://www.radiologyinfo.org>) will insert an intravenous (<http://www.radiologyinfo.org>) catheter (IV line) into a vein in your hand or arm. They will use this IV to inject the contrast material.

You will be placed into the magnet of the MRI unit. The technologist will perform the exam while working at a computer outside of the room. You will be able to talk to the technologist via an intercom.

If your exam uses a contrast material, the technologist will inject it into the intravenous line (IV) after an initial series of scans. They will take more images during or following the injection.

Some MRI exams, such as MR Enterography, may use oral contrast in order to evaluate diseases of the bowel. *For more information, see the MR Enterography page (<https://www.radiologyinfo.org/en/info/mrenterography>)*.

When the exam is complete, the technologist may ask you to wait while the radiologist checks the images in case more are needed.

The technologist will remove your IV line after the exam is over and place a small dressing over the insertion site.

Depending on the type of exam and the equipment used, the entire exam is usually completed in 30 to 50 minutes.

What will I experience during and after the procedure?

Most MRI exams are painless. However, some patients find it uncomfortable to remain still. Others may feel closed-in (claustrophobic) while in the MRI scanner. The scanner can be noisy.

It is normal for the area of your body being imaged to feel slightly warm. If it bothers you, tell the radiologist or technologist. It is important that you remain perfectly still while the images are being taken. This is typically only a few seconds to a few minutes at a

time. You will know when images are being recorded because you will hear and feel loud tapping or thumping sounds. The coils that generate the radio waves make these sounds when they are activated. You will be provided with earplugs or headphones to reduce the noise made by the scanner. You may be able to relax between imaging sequences. However, you will need to keep the same position as much as possible without moving.

You will usually be alone in the exam room. However, the technologist will be able to see, hear, and speak with you at all times using a two-way intercom. They will give you a “squeeze-ball” that alerts the technologist that you need attention right away. Many facilities allow a friend or parent to stay in the room if they have also been screened for safety.

Children will be given appropriately sized earplugs or headphones during the exam. Music may be played through the headphones to help pass the time. MRI scanners are air-conditioned and well-lit.

In some cases, IV injection of contrast material may be given before the images are obtained. The IV needle may cause you some discomfort and you may experience some bruising. There is also a very small chance of skin irritation at the site of the IV tube insertion. Some patients may have a temporary metallic taste in their mouth after the contrast injection.

If you do not require sedation, no recovery period is necessary. You may resume your usual activities and normal diet immediately after the exam. On very rare occasions, a few patients experience side effects from the contrast material. These may include nausea, headache, and pain at the site of injection. It is very rare that patients experience hives, itchy eyes, or other allergic reactions to the contrast material. If you have allergic symptoms, tell the technologist. A radiologist or other doctor will be available for immediate assistance.

Who interprets the results and how do I get them?

A radiologist, a doctor trained to supervise and interpret radiology exams, will analyze the images. The radiologist will send a signed report to your primary care or referring physician, who will share the results with you.

Follow-up exams may be necessary. If so, your doctor will explain why. Sometimes a follow-up exam is done to further evaluate a potential abnormality with additional views or a special imaging technique. Or, a follow-up exam may be to monitor any changes in a known abnormality over time. Follow-up exams are sometimes the best way to see if treatment is working or if a previous finding has changed.

What are the benefits vs. risks?

Benefits

- MRI is a noninvasive imaging technique that does not involve exposure to radiation.
- MR images of soft-tissue structures—such as the liver and many other organs— may be more likely to accurately identify and characterize disease than other imaging methods. This makes MRI an invaluable tool in early diagnosis and evaluation of many focal lesions and tumors.
- MRI has proven valuable in diagnosing a broad range of conditions, including cancer, heart and vascular disease (<http://www.radiologyinfo.org>) , and muscular and bone abnormalities.
- MRI can detect abnormalities that might be obscured by bone with other imaging methods.
- MRI allows physicians to assess the biliary system noninvasively and without contrast injection.
- The MRI gadolinium contrast material is less likely to cause an allergic reaction than the iodine-based contrast materials used for x-rays and CT scanning.
- MRI provides a noninvasive alternative to x-ray, angiography and CT for diagnosing problems of the blood vessels.

Risks

- The MRI exam poses almost no risk to the average patient when appropriate safety guidelines are followed.
- If sedation is used, there is a risk of using too much. However, your vital signs will be monitored to minimize this risk.
- The strong magnetic field is not harmful to you. However, it may cause implanted medical devices to malfunction or distort the images.
- Nephrogenic systemic fibrosis is a recognized complication related to injection of gadolinium (<http://www.radiologyinfo.org>) contrast. It is exceptionally rare with the use of newer gadolinium contrast agents. It usually occurs in patients with serious kidney disease. Your doctor will carefully assess your kidney function before considering a contrast injection.
- There is a very slight risk of an allergic reaction if your exam uses contrast material. Such reactions are usually mild and controlled by medication. If you have an allergic reaction, a doctor will be available for immediate assistance.
- Although there are no known health effects, evidence has shown that very small amounts of gadolinium can remain in the body, particularly the brain, after multiple MRI exams. This is most likely to occur in patients receiving multiple MRI exams over their lifetime for monitoring chronic or high-risk health conditions. The contrast agent is mostly eliminated from the body through the kidneys. If you are a patient in this category, consult with your doctor about the possibility of gadolinium retention, as this effect varies from patient to patient.
- Manufacturers of intravenous contrast indicate mothers should not breastfeed their babies for 24-48 hours after contrast medium is given. However, both the American College of Radiology (ACR) and the European Society of Urogenital Radiology note that the available data suggest that it is safe to continue breastfeeding after receiving intravenous contrast. *For further information please consult the ACR Manual on Contrast Media (<https://www.acr.org/Clinical-Resources/Contrast-Manual>) and its references.*

What are the limitations of abdominal and pelvic MRI?

High-quality images depend on your ability to remain perfectly still and follow breath-holding instructions while the images are being recorded. If you are anxious, confused or in severe pain, you may find it difficult to lie still during imaging.

A person who is very large may not fit into the opening of certain types of MRI machines.

Implants and other metallic objects can make it difficult to obtain clear images. Patient movement can have the same effect.

Breathing may cause artifacts (<http://www.radiologyinfo.org>) , or image distortions, during MRI. Bowel motion may also cause artifacts. This is less of a problem with state-of-the art scanners and techniques.

Present data show no convincing evidence that non contrast MRI harms the fetus of a pregnant woman. However, if the need for the exam is not time sensitive your doctor may delay the exam until after delivery. MRI gadolinium contrast agents are generally avoided during pregnancy except in very specific circumstances. Your doctor will discuss the benefits and risks of any MRI procedure with you. Doctors may perform MRI after the first trimester to assess the fetus for findings that are not fully evaluated by ultrasound.

MRI may not always distinguish between cancer tissue and benign tumors or other conditions, such as edema (<http://www.radiologyinfo.org>) .

An MRI exam typically costs more and may take more time than other imaging exams. Talk to your insurance provider if you have concerns about the cost of MRI.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.