

Ataxia-Child

Not being able to move in a coordinated fashion is referred to as ataxia. Signs of ataxia include clumsiness, unsteady gait, nystagmus (rapid, uncontrolled eye movements), and dysdiadochokinesis (inability to perform rapid alternating movements like foot tapping).

Ataxia may be genetic or acquired, such as from illness or trauma. Acute-onset ataxia develops within hours or days. In children with acute ataxia and no recent history of trauma, MRI head (<https://www.radiologyinfo.org/en/info/mri-brain>) without and with contrast, MRI head without contrast, or CT head (<https://www.radiologyinfo.org/en/info/headct>) without contrast is usually appropriate. MR angiography (MRA) (<https://www.radiologyinfo.org/en/info/angiomr>) head and/or neck, MRI spine (<https://www.radiologyinfo.org/en/info/spinemr>), and CT angiography (CTA) (<https://www.radiologyinfo.org/en/info/angiocr>) head and/or neck may be appropriate.

For acute ataxia (after traumatic head or neck injury), MRI head without contrast or CT head without contrast is usually appropriate. MRA head and/or neck and CTA head and/or neck may be appropriate.

For recurrent ataxia (episodes of ataxia followed by relatively symptom-free periods), MRI head without contrast is usually appropriate. MRI spectroscopy head or MRI head without and with contrast may be appropriate.

For chronic progressive ataxia (symptoms typically appear and increase in severity over at least 2 months), MRI head without and with contrast or MRI head without contrast is usually appropriate. MRI complete spine without or with/without contrast and CT head without contrast may be appropriate.

For chronic nonprogressive ataxia (symptoms typically appear in the first year of life, last more than 2 months, and do not change in severity), MRI head without contrast is usually appropriate. MRI complete spine without contrast and CT head without contrast may be appropriate.

— By Emily Guernsey and Orit A. Glenn, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or

the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2025 Radiological Society of North America, Inc.