

## Cervical Neck Pain or Cervical Radiculopathy

Appropriate cervical spine (c-spine) imaging and use of intravenous contrast (venous dye) for neck pain varies depending on clinical scenario.

In individuals with new or increasing nontraumatic neck pain (no high-risk factors), x-rays (<https://www.radiologyinfo.org/en/info/bonerad>) are usually appropriate as first imaging test; MRI (<https://www.radiologyinfo.org/en/info/headmr>) or CT (<https://www.radiologyinfo.org/en/info/headct>) without contrast may be appropriate.

In cases with spinal nerve irritation (pinched nerve), MRI without contrast is usually appropriate, and x-rays or CT without contrast may be appropriate.

If there is history of prior c-spine surgery, x-rays or CT without contrast is usually appropriate; MRI with or without contrast (or both) or CT myelography (<https://www.radiologyinfo.org/en/info/myelography>) (contrast injection in space around spinal cord) may be appropriate.

If infection is suspected, MRI without and with contrast is usually appropriate; x-rays, CT, or MRI with or without contrast may be appropriate.

In individuals with cancerous tumors, MRI without and with or only without contrast is usually appropriate; x-rays, CT with or without contrast, MRI with contrast, or bone scan or CT may also be appropriate.

With headaches (<https://www.radiologyinfo.org/en/info/headache>) originating in the c-spine without weakened nerve function, MRI or CT without contrast may be appropriate, as well as x-rays or nerve block (<https://www.radiologyinfo.org/en/info/nerveblock>) injection.

In chronic neck pain, x-rays are usually appropriate as initial imaging. MRI without contrast may be appropriate.

If x-rays show degenerative changes, MRI without contrast is usually appropriate. CT without contrast or CT myelography may be appropriate.

If x-rays show hardening of the connecting fibers of the neck vertebrae, CT without contrast is usually appropriate; CT myelography or MRI without contrast may be appropriate.

— By Susan Anemone and Bruno Policeni, MD, MBA. This information originally appeared in the *Journal of the American College of Radiology*.

### Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

## Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2023 Radiological Society of North America, Inc.