

Chronic Ankle Pain

When ankle pain continues for 6 weeks or more, it is considered chronic. The most appropriate first imaging test is an x-ray (<https://www.radiologyinfo.org/en/info/bonerad>) . X-rays provide general information about bone and tissue abnormalities. It is almost always the only imaging test of the ankle needed.

In people who have ankle arthritis (<https://www.radiologyinfo.org/en/info/arthriti>) and have pain that requires treatment, it may be appropriate to figure out which of the joints in the ankle is causing the pain. Either CT (<https://www.radiologyinfo.org/en/info/bodyct>) or MRI (<https://www.radiologyinfo.org/en/info/muscmr>) without contrast is appropriate to locate the joint causing the pain. Using imaging to guide a needle to inject anesthetic in the area of the pain to help confirm the cause is usually appropriate, too.

Some chronic ankle conditions cannot be seen on x-ray. When these conditions are suspected, MRI without intravenous contrast is appropriate. CT or MR arthrography (<https://www.radiologyinfo.org/en/info/arthrog>) may also be appropriate, in which case an x-ray dye is injected before the test is performed. These conditions include:

- Osteochondral lesion: an injury affecting the talus bone (a supporting bone in the ankle) and the surrounding cartilage. CT or MRI may be appropriate.
- Tendon abnormality: inflammation of the tendon, commonly known as tendonitis. Ultrasound (<https://www.radiologyinfo.org/en/info/musculos>) or MRI may be appropriate.
- Ankle instability: residual wear and tear on the ligaments makes the ankle unstable. Ultrasound or MRI may be appropriate.
- Ankle impingement syndrome: overuse and microtears of the tendon causing pain and limiting ankle range of motion. MRI, CT or ultrasound may be appropriate.
- No obvious suspected disorder: MRI, CT, ultrasound, or image-guided local anesthetic injections may be appropriate.

— By Frank J. Rybicki Jr. and MacArinze Ojiaku, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple

distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2025 Radiological Society of North America, Inc.