

Chronic Liver Disease

Chronic liver disease can be caused by a variety of conditions but is most commonly caused by nonalcoholic fatty liver disease (<https://www.radiologyinfo.org/en/info/fatty-liver-disease>), hepatitis C, hepatitis B, and alcohol-related liver disease. Over time, these conditions can lead to fibrosis (thickening, scarring of tissue) and to cirrhosis, in which the damage is so extensive that the liver does not function normally. These diseases can also result in liver cancer (hepatocellular carcinoma [HCC]).

In individuals with chronic liver disease, imaging tests are useful to confirm the presence and severity of fibrosis. MR elastography abdomen finds stiffening of the liver, and ultrasound shear wave elastography abdomen is usually appropriate for diagnosing and staging of fibrosis.

Individuals who have chronic viral hepatitis and cirrhosis are at high risk for developing HCC and should undergo screening with imaging. Ultrasound abdomen (<https://www.radiologyinfo.org/en/info/abdominus>), MRI abdomen (<https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis>) without and with intravenous (IV) contrast, and MRI abdomen without and with hepatobiliary contrast are usually appropriate imaging tests to diagnose HCC. MRI abdomen without IV contrast may also be appropriate.

For individuals with HCC, surveillance after treatment with imaging tests is recommended to detect a possible return of cancer. MRI abdomen without and with IV contrast, CT abdomen with IV contrast multiphase, CT abdomen without and with IV contrast, and MRI abdomen without and with hepatobiliary contrast are usually appropriate tests. The current guidelines recommend a CT or MRI scan every 3 to 6 months for 2 years and then every 6 to 12 months after the cancer is removed.

— By Emmanuel Budis and Jennifer W. Uyeda, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.