

Crohn Disease—Child

Crohn disease (<https://www.radiologyinfo.org/en/info/crohns-disease>) is a common inflammatory bowel disease that can affect children, adolescents, and adults. It is diagnosed using a combination of physical examination, blood work, endoscopy, and imaging tests.

Symptoms of Crohn disease can include weight loss, loose stools, vomiting, and intermittent abdominal pain. In children with suspected Crohn disease, *CT enterography (CTE)* (<https://www.radiologyinfo.org/en/info/ctenterography>), *MR enterography (MRE)* (<https://www.radiologyinfo.org/en/info/mreenterography>), or *MRI of the abdomen/pelvis* (<https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis>) without and with intravenous contrast is usually appropriate. *Ultrasound (US) of the abdomen* (<https://www.radiologyinfo.org/en/info/abdominus>), *CT abdomen/pelvis* (<https://www.radiologyinfo.org/en/info/abdominct>) with contrast, fluoroscopic *upper gastrointestinal series* (<https://www.radiologyinfo.org/en/info/uppergi>) with small bowel follow-through (UGI-SBFT), or *MRI abdomen/pelvis* without contrast may be appropriate. For enterography studies, children drink contrast before imaging to stretch the bowel for optimal imaging.

The severity of Crohn disease symptoms varies over time. For children with known Crohn disease with worsening symptoms, *CT abdomen/pelvis* with contrast, *CTE*, *MRE*, or *MRI abdomen/pelvis* without and with contrast is usually appropriate. *US abdomen, abdomen x-ray* (<https://www.radiologyinfo.org/en/info/abdominrad>), fluoroscopic *UGI-SBFT*, or *MRI abdomen/pelvis* without contrast may also be appropriate.

For evaluation of disease surveillance or monitoring therapy, *CTE*, *MRE*, or *MRI abdomen/pelvis* without and with contrast is usually appropriate. *US abdomen, US abdomen with microbubble contrast, CT abdomen/pelvis* with contrast, fluoroscopic *UGI-SBFT*, *MRI abdomen/pelvis* without contrast, or *fluorine-18-2-fluoro-2-deoxy-D-glucose PET/CT* (<https://www.radiologyinfo.org/en/info/pet>) may also be appropriate.

If complications develop around the anus such as fistula or infected fluid collections, *MRI pelvis* with contrast or *MRI pelvis* without and with contrast is usually appropriate. *Transperineal ultrasound of the pelvis, CT pelvis* with contrast, or *MRI pelvis* without contrast may also be appropriate.

For more information, see the Crohn disease (<https://www.radiologyinfo.org/en/info/crohns-disease>) *page.*

— By Emily K. Guemsey, BS and Aya Kamaya, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to

other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.