

Lower Extremity Arterial Revascularization - Post-Therapy Imaging

The goal of endovascular angioplasty (<https://www.radiologyinfo.org/en/info/angioplasty>) and of surgical bypass is to restore normal blood flow to the diseased arteries in the legs for individuals with peripheral artery disease. Both procedures need imaging follow-up to check for lesions, typically narrowing of arteries that may come back and cause symptoms. Often, noninvasive testing is done first to help choose the appropriate imaging test. The first test is usually measurement of the ankle-brachial index (ABI), a noninvasive test in which blood pressure cuffs are placed on the legs to measure local pressures. Other noninvasive tests can also be used.

For individuals without symptoms after their procedure, it is appropriate to use a baseline ultrasound (US) (<https://www.radiologyinfo.org/en/info/vascularus>) examination in addition to ABI and other noninvasive tests. For patients with symptoms such as severe leg cramps during exercise, pain at rest, and wounds that do not heal properly after the procedure, imaging can include CT (<https://www.radiologyinfo.org/en/info/angiocr>) or MR angiography (<https://www.radiologyinfo.org/en/info/angiomr>) in addition to US. If the symptoms are more severe, including cold and discolored limbs with a very low pulse, the primary object is to save the leg from amputation. In extremely severe cases, it is appropriate to skip imaging and go directly to surgery. In nonthreatening but severe cases, arteriography (x-rays after a dye is injected into blood vessels through a catheter) is the most appropriate follow-up. Arteriography can provide images of lesions that limit flow and cause drops in leg blood pressure. In less severe cases, both US and CT angiography are appropriate imaging tests. *For more information, see the Peripheral Artery Disease (<https://www.radiologyinfo.org/en/info/pad>) page.*

— By Frank Rybicki Jr., Karin E. Dill, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2023 Radiological Society of North America, Inc.