

Management of Liver Cancer

Treatment options for liver cancer depend on the number, sizes, and locations of tumors.

Hepatocellular Cancer (Most Common)

- For a single tumor <3 cm in a cirrhotic liver, liver transplantation, percutaneous ablation (<https://www.radiologyinfo.org/en/info/thermal-ablation-therapy>) (destroying tumor using microwave or thermal energy), or liver resection (piece of liver removed) is usually appropriate.
- For a single tumor 3 to 5 cm in a cirrhotic liver, liver transplantation, combination therapies, liver resection, and treatments to block tumor blood flow are usually appropriate.
- For multiple tumors in a cirrhotic liver (both lobes), with at least 1 tumor >5 cm, blocking tumor blood flow and systematic treatments (chemotherapy) are usually appropriate.
- For single or multiple tumors in a cirrhotic liver with vascular invasion (cancer has grown into a blood vessel), systematic therapies and transarterial radioembolization (<https://www.radiologyinfo.org/en/info/radioembol>) (blocking tumor blood flow using radio waves) are usually appropriate.

Cholangiocarcinoma (Bile Duct Cancer)

- For bile duct cancer in peripheral liver <3 cm, with no biliary ductal dilatation, vascular invasion, regional lymphadenopathy, or metastases, liver resection and percutaneous ablation liver are usually appropriate.
- For cancer in central bile ducts >3 cm with poorly defined margins, vascular invasion, and periportal lymphadenopathy, systematic therapies are usually appropriate.

Metastatic Liver Disease (Cancer Has Spread to the Liver From Elsewhere in Body)

- For a multifocal metastatic neuroendocrine tumor, long-acting somatostatin analogs (which slow down production of hormones), peptide receptor radionuclide therapy (which delivers radiation directly to cancer), and blocking blood flow to the tumor are usually appropriate.
- For single metastases to the liver from colorectal cancer, systematic therapies, liver resection, and percutaneous ablation liver are usually appropriate.
- For multifocal metastases to both the right and left liver from colorectal carcinoma: systematic therapies are usually appropriate.

— By Samantha Greben and Aya Kamaya, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR

and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.