

Monitoring Response to Neoadjuvant Systemic Therapy for Breast Cancer

Neoadjuvant chemotherapy (NAC) is treatment given before surgery to help shrink tumors and reduce the chances of the cancer spreading (metastasis). The tests and procedures discussed here apply to adult female, male, transfeminine (male-to-female), and transmasculine (female-to-male) patients with known breast cancer.

Before NAC, ultrasound (US) breast (<http://www.radiologyinfo.org/breastus>) , digital breast tomosynthesis (DBT) (<http://www.radiologyinfo.org/tomosynthesis>) diagnostic, mammography (<http://www.radiologyinfo.org/mammo>) diagnostic, and MRI breast (<http://www.radiologyinfo.org/breastmr>) without and with intravenous (IV) contrast are usually appropriate.

After initiation or completion of NAC, US breast, DBT diagnostic, mammography diagnostic, and MRI breast without and with IV contrast are usually appropriate.

For people with known clinically node-negative breast cancer (<http://www.radiologyinfo.org/breast-cancer>) (cancer has not spread to the lymph nodes), US axilla (US underarm or armpit) is usually appropriate before NAC.

For people with known clinically node-positive breast cancer (cancer has spread to the lymph nodes), US axilla is usually appropriate. MRI breast without and with IV contrast and fluorine-18-2-fluoro-2-deoxy-D-glucose (FDG)-PET/CT (<http://www.radiologyinfo.org/pet>) skull base to mid-thigh may be appropriate.

After NAC completion, for people with known clinically node-negative breast cancer, there are no imaging tests recommended.

For assessment of response to NAC, for people with known breast cancer and suspicion of metastasis, bone scan (<http://www.radiologyinfo.org/bone-scan>) whole body, CT chest abdomen pelvis (<http://www.radiologyinfo.org/abdominect>) with IV contrast, and FDG-PET/CT skull base to mid-thigh are usually appropriate.

For people with known axillary lymph node-positive breast cancer, US axilla are usually appropriate after NAC.

For people with known breast cancer and axillary imaging suspicious for metastasis, US-guided core biopsy (<http://www.radiologyinfo.org/breastbius>) axillary node and US-guided fine needle aspiration biopsy axillary node are usually appropriate.

For more information, see the Breast Cancer page (<http://www.radiologyinfo.org/breast-cancer>) .

—By Rachael Newman and Sherry S. Wang, MBBS. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information

is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2025 Radiological Society of North America, Inc.