

Noninvasive Clinical Staging of Primary Lung Cancer

Lung cancer (https://www.radiologyinfo.org/en/info/lung-cancer) is the leading cause of cancer-related deaths in both men and women in the United States. The major cause of lung cancer is cigarette smoking, but other risks include exposure to secondhand smoke, environmental radon, occupational exposures, and air pollution.

There are two main forms of lung cancer: non–small-cell lung cancer (NSCLC) and small-cell lung cancer (SCLC). NSCLC is the most common cancer and accounts for approximately 85% of cases, and SCLC accounts for approximately 15%. Staging determines the extent that the cancer has grown and spread. Numbers from 0 to 4 (I, II, III, IV) refer to how far the cancer has spread, with stage IV being the most advanced stage.

Usually, appropriate imaging tests for the staging of NSCLC are an initial CT chest (https://www.radiologyinfo.org/en/info/chestct) with intravenous (IV) contrast or without IV contrast followed by FDG-PET/CT (https://www.radiologyinfo.org/en/info/pet) skull base to mid-thigh (uses IV radioactive sugar to find metabolic activity in cancer cells) and MRI head (https://www.radiologyinfo.org/en/info/mri-brain) without/with IV contrast (to look for cancer spread to the brain) which are obtained based on the initial CT chest and other test results. CT head (https://www.radiologyinfo.org/en/info/headct), CT abdomen/pelvis (https://www.radiologyinfo.org/en/info/abdominct), MRI chest (https://www.radiologyinfo.org/en/info/chestmr), MRI abdomen (https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis), and whole-body bone scan (https://www.radiologyinfo.org/en/info/bone-scan) may be appropriate depending on the situation. Similarly, to stage SCLC, CT chest with IV contrast, FDG-PET/CT skull base to mid-thigh, MRI head without/with IV contrast, and CT abdomen/pelvis with IV contrast are usually appropriate. CT head, CT abdomen/pelvis, MRI chest, MRI abdomen, and whole-body bone scan may be appropriate.

For more information, visit the Lung Cancer (https://www.radiologyinfo.org/en/info/lung-cancer) page.

— By Elle Sutherland and Gregory J. Czuczman, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (http://www.radiologyinfo.org) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at http://www.radiologyinfo.org to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright ® 2024 Radiological Society of North America, Inc.	
vasive Clinical Staging of Primary Lung Cancer ight© 2024, RadiologyInfo.org	Page 2 of 2 Reviewed Jan-24-2024