

Pancreatic Cyst

Incidental pancreatic cysts are fluid-filled pockets that develop in or on the pancreas, typically identified on imaging tests done for another purpose. Common pancreatic cysts include intraductal papillary mucinous neoplasms, serous cystadenomas, and mucinous cystic neoplasms.

For incidental pancreatic cysts less than or equal to 2.5 cm in size, MRI abdomen (<https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis>) without and with contrast with magnetic resonance cholangiopancreatography (MRCP) (<https://www.radiologyinfo.org/en/info/mrcp>) (MRI test designed to assess the pancreatic system) is usually appropriate. Multiphase CT abdomen (<https://www.radiologyinfo.org/en/info/abdominct>) (images taken at different time points) and MRI abdomen without contrast with MRCP may be appropriate.

For incidental cysts greater than 2.5 cm with no additional worrisome features, MRI abdomen without and with contrast with MRCP is usually appropriate. Multiphase CT abdomen with contrast, MRI abdomen without contrast with MRCP, and ultrasound abdomen endoscopic may be appropriate.

For cysts greater than 2.5 cm with worrisome features, abdominal endoscopic ultrasound (<https://www.radiologyinfo.org/en/info/abdominus>) or MRI abdomen without and with contrast with MRCP is usually appropriate. Multiphase CT abdomen with contrast and MRI abdomen without contrast with MRCP may be appropriate.

If the pancreatic duct (tube for pancreatic juices for digestion) is greater than 7 mm, abdominal endoscopic ultrasound, MRI abdomen without and with contrast with MRCP, and MRI abdomen without contrast with MRCP are usually appropriate. Multiphase CT abdomen with contrast may be appropriate.

Multiphase CT abdomen with contrast, MRI abdomen without and with contrast with MRCP, and MRI abdomen without contrast with MRCP are usually appropriate for follow-up imaging.

— By Rachel Newman and Samantha Lynn Heller, PhD, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or

the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.