

## **Pelvic Floor Dysfunction in Female Patients**

Pelvic floor dysfunction is a group of conditions that is common in women, especially after menopause. It is caused by weakness of the muscles in the pelvis. This weakness can affect urination, defecation (bowel movements), and support of pelvic organs, leading to prolapse (bulging out) of the vagina or rectum.

These conditions can cause a lot of discomfort. Accurate diagnosis is important for proper treatment.

Imaging tests can help diagnose the problem. These include fluoroscopy (video x-rays), MRI, and ultrasound (US). These tests may require the use of contrast (liquid or dye) that is given orally, injected in the vein, or placed in the vagina, bladder, or rectum, depending on the test. Some tests are done while the patient is urinating or defecating.

The usually appropriate imaging tests are based on the specific type of pelvic floor problem. Fluoroscopy and MRI are usually appropriate tests for women with prolapse (bulging out) of the vagina or rectum.

Fluoroscopy is usually appropriate in the initial diagnosis of people with urination problems (urine leaking, urinating frequently, strong urge to urinate, and difficulty emptying the bladder). Specialized types of US and MRI may also be appropriate.

In people with defecation problems (incontinence, straining, strong urge to defecate, difficulty passing bowel movements), specialized types of fluoroscopy, US, and MRI are usually appropriate.

In people who have had pelvic floor surgery, or with other chronic complications, specialized MRI tests are usually appropriate.

For more information, see the MR Defecography (http://www.radiologyinfo.orgdefecography-mri) page.

— By Anika Shah and Nina S. Vincoff, MD. This information originally appeared in the *Journal of the American College of Radiology*.

## Disclaimer

This information is copied from the RadiologyInfo Web site (http://www.radiologyinfo.org) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at http://www.radiologyinfo.org to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

## Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright ® 2025 Radiological Society of North America, Inc.	
ic Floor Dysfunction in Female Patients /right© 2025, RadiologyInfo.org	Page 2 of Reviewed Sep-23-202