

Radiologic Management of Uterine Leiomyomas

Leiomyomas (fibroids) in the uterus are benign tumors and are common in women of reproductive age. They are treated when they cause abnormal bleeding, pain, pressure, or bloating.

The usually appropriate treatments are (1) oral medication; (2) uterine artery embolization (UAE) (<https://www.radiologyinfo.org/en/info/uterine-fibroid-treatment>) , or blocking the arteries that feed the fibroids; (3) myomectomy, or the surgical removal of the fibroids; and (4) hysterectomy, or surgical removal of the entire uterus. High-frequency focused ultrasound (HIFU) ablation surgery, in which sound waves are used to break down the fibroid, can also be used.

In middle-aged woman with multiple fibroids, excessive menstrual bleeding, urinary frequency, and bloating who are not pregnant and do not want a future pregnancy, UAE or hysterectomy is usually appropriate. If there is uterine wall fibroid and growth of uterine lining tissue into the uterine wall, surgical removal of uterine lining may be appropriate.

In middle-aged woman with pelvic discomfort and uterine lining fibroid, UAE, myomectomy, or hysterectomy is usually appropriate. Oral medication may also be appropriate. If there is fibroid compressing the rectum, UAE, myomectomy, or hysterectomy is usually appropriate.

In childbearing-age woman who do not want a future pregnancy but are concerned about losing femininity after hysterectomy, UAE is usually appropriate. Hysterectomy may be appropriate but would result in complete loss of fertility.

In childbearing-age women with multiple fibroids in the uterine wall who may want a future pregnancy, UAE or myomectomy is usually appropriate. Ultrasound ablation (HIFU) may also be appropriate.

— By Susan Anemone and MacArinze Ojiaku, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

