

Sinonasal Disease

Sinonasal disease includes rhinosinusitis (inflammation of the nose or sinuses). “Acute” means that symptoms last less than 4 weeks, and “chronic” means that symptoms last more than 12 weeks.

For acute rhinosinusitis alone, no imaging tests are recommended.

For acute rhinosinusitis with possible orbit (eye) or brain involvement, head MRI (<https://www.radiologyinfo.org/en/info/mri-brain>) or orbits, face, and neck MRI without and with contrast or maxillofacial CT with contrast is usually appropriate. Head MRI without contrast or head CT with contrast may be appropriate.

For people who may be candidates for surgery, maxillofacial CT without contrast is usually appropriate. Orbita, face, and neck MRI or CT of the sinuses (<https://www.radiologyinfo.org/en/info/sinusct>) (<https://www.radiologyinfo.org/en/info/sinusct>) without contrast or maxillofacial CT with contrast may be appropriate.

For acute sinusitis that quickly gets worse or fungal sinusitis that may be spreading, orbits, face, and neck MRI without and with contrast, maxillofacial CT with contrast, or maxillofacial CT without contrast is usually appropriate. Head MRI without and with contrast, head MRI, or orbits, face, and neck MRI without contrast or head CT with contrast may be appropriate.

For possible sinonasal mass, orbits, face, and neck MRI without and with contrast, maxillofacial CT with contrast, or maxillofacial CT without contrast is usually appropriate. Head MRI without and with contrast, head MRI, or orbits, face, and neck MRI without contrast or head CT with contrast may be appropriate.

For possible leaking of fluid surrounding the brain and spinal cord, maxillofacial CT without contrast is usually appropriate. Head MRI or orbits, face, and neck MRI without contrast; head CT cisternography or DTPA cisternography; or sinus single-photon emission computed tomography (<https://www.radiologyinfo.org/en/info/pet>) or single-photon emission computed tomography/CT may be appropriate. These tests determine whether there is fluid in the sinuses.

— By Sandra V. Kotsis, MPH and Aya Kamaya, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.