

## **Suspected Pulmonary Embolism**

Imaging tests are not necessary for many patients suspected of having a blood clot in the lungs known as a pulmonary embolism (PE) (https://www.radiologyinfo.org/en/info/pulmonary-embolism). For medically stable patients, the risk of a PE should be assessed first by asking standard questions about the patient and his or her symptoms. If the answers to the questions result in a low risk score, no further testing is required.

If the answers indicate the possibility of PE, a blood test (D-dimer) to check for a substance released when a blood clot (https://www.radiologyinfo.org/en/info/bloodclot) breaks up is recommended. If the test comes back negative, no further testing is required. The D-dimer test should not be used for anybody expected to have blood clots due to other things, such as recent surgery or trauma, or for pregnant women.

If the answers to the standard questions indicate a high risk of PE and the D-dimer test is positive, in most cases a pulmonary CT angiography (CTA) (https://www.radiologyinfo.org/en/info/angioct) —a CT scan to look at the blood vessels in the lungs—is the next step. For people with symptoms of a blood clot in the lower legs, especially for pregnant women, an ultrasound (https://www.radiologyinfo.org/en/info/venousus) Doppler of the legs is often the first choice to reduce radiation exposure. Eighty percent of PEs are associated with blood clots in the lower legs. In addition, a chest x-ray (https://www.radiologyinfo.org/en/info/chestrad) may be performed to rule out other causes such as pneumonia or fluid in the lungs. Ventilation and perfusion nuclear medicine scans are sometimes used in place of CTA.

For more information, visit the Pulmonary Embolism (https://www.radiologyinfo.org/en/info/pulmonary-embolism) page.

— By Andrea Borondy Kitts, MPH. This information originally appeared in the *Journal of the American College of Radiology*.

## Disclaimer

This information is copied from the RadiologyInfo Web site (http://www.radiologyinfo.org) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at http://www.radiologyinfo.org to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

## Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright ® 2024 Radiological Society of North America, Inc.