



Dementia

Dementia refers to a group of symptoms that indicate brain function has been impaired. The condition is characterized by a decline in language, learning and/or memory that is severe enough to interfere with daily function and independence.

A diagnosis of dementia is made via a variety of tests to rule out other, treatable conditions and may include physical and neurological examinations as well as cognitive testing. Some dementias are incurable while the progression of other dementias may be slowed or reversed with treatment.



What is dementia?

Dementia refers to a group of symptoms that indicate brain function has been impaired, usually by disease or injury. It is characterized by a decline in one or more areas of conscious mental activity—such as language, learning and memory—severe enough to interfere with daily function and independence. Symptoms may include:

- memory loss
- language problems
- inability to focus
- depression
- difficulty with routine daily activities, such as eating or getting dressed
- difficulty controlling emotions
- changes in personality
- agitation

Dementia affects an estimated 47.5 million people worldwide; this number is expected to grow significantly as the population ages.

Many different diseases can cause dementia. Alzheimer's disease is the leading cause, accounting for up to one half to two-thirds of cases. The second most common cause is vascular dementia, also known as

vascular cognitive impairment (VCI), which often occurs after a stroke when blood flow to certain areas of the brain is blocked. Mixed dementia, where a person has more than one type of dementia, is often found in older people. However, dementia is not a normal part of aging.

How is dementia diagnosed and evaluated?

Dementia diagnosis uses a variety of tests to rule out other, treatable conditions such as depression, vitamin B12 deficiency, hydrocephalus and hypothyroidism. Physical and neurological examinations and cognitive testing are usually part of the analysis.

Brain imaging is a useful complement to laboratory tests to rule out other treatable conditions such as brain tumor. Your doctor may order one of the following imaging tests:

MRI of the head uses a powerful magnetic field, radio frequency pulses and a computer to produce detailed pictures of the brain to detect brain abnormalities and diagnose conditions in the brain. Sometimes contrast material is injected into an arm vein to enhance the images. Head MRI may also reveal whether a stroke has recently occurred. See the Safety page for more information about MRI. In many cases the physicians may order a second MRI to assess changes that have occurred in the interval.

Brain functional magnetic resonance imaging (fMRI) uses MR imaging to measure the tiny metabolic changes that take place in an active part of the brain. fMRI is used to help assess how brain function has been impacted by stroke, trauma or degenerative disease such as Alzheimer's.

CT of the head combines special x-ray equipment with sophisticated computers to produce multiple images or pictures of the inside of the head. CT can detect atrophy related to dementia and provide detailed information on head injuries, stroke, and other brain diseases. Sometimes contrast material is injected into an arm vein to enhance the images. See the Safety page for more information about CT.

PET is a diagnostic examination that uses a small amount of radioactive material (called a radiotracer) to help diagnose and determine the severity of a variety of diseases. A combined PET/CT exam fuses images from a PET and CT scan together to provide detail on both the anatomy (from the CT scan) and function (from the PET scan) of the brain. A special form of PET can help identify abnormal clusters of proteins associated with Alzheimer's disease.

Lumbar puncture, also known as a spinal tap, is a diagnostic test that uses a needle to remove a small amount of cerebrospinal fluid—the fluid that surrounds the brain and spinal cord. Research suggests that people with early-stage Alzheimer's disease may have abnormal protein clusters in their cerebrospinal fluid.

If neurologists suspect a treatable dementia called normal pressure hydrocephalus (NPH), they may order a test called radionuclide cisternography. In this test a radiotracer is injected via lumbar puncture into the cerebrospinal fluid. The pattern of distribution of this tracer over the brain with time helps diagnose normal-pressure hydrocephalus.

How is dementia treated?

While there is no cure for some dementias, the progression of some types of dementia can be slowed or even reversed with treatment. Options include:

- Treating the cause of dementia when there is a treatable cause: This includes hormonal treatment for hypothyroidism, treating hydrocephalus with shunting, evacuation of subdural collections, etc.
- Cholinesterase inhibitors, a type of medication that may slow the progress of Alzheimer's disease by helping people improve their attention and working memory.
- Medications to manage blood pressure, cholesterol, triglycerides, diabetes, and problems with blood clotting.
- Procedures to improve blood flow to the brain, such as carotid endarterectomy or Angioplasty and Vascular Stenting.
- Lifestyle modifications, such as following a healthy diet and adding an exercise regimen, quitting smoking and quitting or decreasing alcohol consumption.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2017 Radiological Society of North America, Inc.