



## Osteoporosis

Osteoporosis is a disease characterized by thinning and deterioration of bone tissue and loss of density. The condition can progress over time and cause bone to become increasingly porous and weak and break more easily.

Your doctor may use x-ray, body CT, spine CT or a bone density (DEXA) scan to help diagnose your condition and assess your risk for fracture. Osteoporosis may cause vertebral compression fractures in the spine; if painful, these fractures may be treated with vertebroplasty or kyphoplasty.



## What is osteoporosis?

Osteoporosis is a disease characterized by thinning and deterioration of bone tissue with loss of calcification and density. Osteoporosis can progress over time. This can cause the bone to become fragile, increasingly porous (as the name of the disease would suggest), and to break more easily. When viewed microscopically, healthy bone looks somewhat like a honeycomb. In cases where osteoporosis is present, the holes and spaces in the honeycomb appear much bigger than those found in a healthy bone.

Women and older adults are more at risk for developing osteoporosis. Heredity, low body weight, and chronic use of certain medications (such as steroids) are also risk factors for the disease.

Lifting, bending, bumping into furniture and even sneezing can cause a bone to break in people affected by osteoporosis. Fractures of the hip, spine or wrist are most common, but other bones are also susceptible to breaks.

Osteoporosis can be present for years without any noticeable symptoms, but signs can include:

- Severe back pain
- Loss of height over time
- A stooped posture
- Bone fractures from minor injury

## How is osteoporosis diagnosed and evaluated?

To diagnose osteoporosis and assess your risk of fracture and determine your need for treatment, your doctor will most likely order a bone density scan.

This exam is used to measure bone mineral density (BMD). It is most commonly performed using dual-energy x-ray absorptiometry (DXA or DEXA) or bone densitometry. The amount of x-rays absorbed by tissues and bone is measured by the DXA machine and correlates with bone mineral density.

The DXA machine converts raw density information to your T score and Z score. The T score measures the amount of bone you have in comparison to a normal population of younger people and is used to estimate your risk of developing a fracture. Your Z score measures the amount of bone you have in comparison to those in your age group. This number can help indicate whether there is a need for further medical tests.

The following procedures can be performed to determine bone injury or fractures due to osteoporosis:

- **Bone x-ray:** Bone x-ray produces images of bones within the body, including the hand, wrist, arm, elbow, shoulder, foot, ankle, leg (shin), knee, thigh, hip, pelvis or spine. It aids in the diagnosis of fractured bones, which are sometimes a result of osteoporosis.
- **CT scan of the spine:** CT scanning of the spine is performed to assess for alignment and fractures. It can be used to subjectively measure bone density and determine whether vertebral fractures are likely to occur. This technique is called quantitative CT (QCT).
- **MRI of the spine:** Magnetic resonance imaging of the spine is performed to evaluate vertebral fractures for evidence of underlying disease, such as cancer, and to assess the newness of the fracture. New fractures demonstrate a better response to treatment by vertebroplasty and kyphoplasty in certain clinical situations.

## How might osteoporosis be treated?

Compounds called bisphosphonates are the standard of care for the treatment of osteoporosis. A prescription is required for this medication and medical evaluation is required before treatment.

Compression fractures in the vertebra can occur as a result of osteoporosis. In these cases, vertebroplasty and kyphoplasty, performed by a radiologist, may be an option to treat painful spine fractures. With vertebroplasty, image guidance is used to inject a special cement mixture through a hollow needle into the fractured bone. In kyphoplasty, a balloon is inserted through the needle into the fractured bone to create a cavity. Once the balloon is removed, a cement mixture is injected into the cavity.

In some cases of compression fracture, surgical treatment by a neurosurgeon may be required, especially if there is evidence of severe narrowing of your spinal canal.

## Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

## Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2018 Radiological Society of North America, Inc.