

How much dose do I get from different imaging procedures?

When it comes to radiation dose, all imaging procedures are not the same. Some procedures, like ultrasound and magnetic resonance imaging (MRI), use no x-rays or nuclear radiation. Procedures that use x-rays (like standard x-ray or computed tomography) or radioactive materials (nuclear medicine) vary widely in dose. Dose depends on the type of procedure and the part of the body being examined.

Potential risk can be expressed in different ways. Radiation dose can be compared to levels from naturally existing radiation in the environment, or a qualitative assessment of risk can be given.

Effective dose



Ankle x-ray

Effective dose is the dose quantity that is used to assess risks from diagnostic medical imaging procedures. It is expressed in millisieverts (mSv).

Effective dose takes many factors into account, including the sensitivity of different body organs and tissues.

For example, an ankle x-ray has a different effective dose than a coronary angiogram.


However, effective dose is not intended to apply to a specific patient. The actual risk to a patient might be higher or lower, depending on the size of the patient and the type of procedure.

Natural background radiation in the United States

We are always exposed to background radiation. It comes from the air, sky, ground and the food we eat. It is natural to our environment.

To put dose from medical imaging in perspective, we compare imaging dose to the time it takes to reach the same dose from natural background radiation.

Over one year's time, our dose from natural background radiation is approximately 3 mSv.



Qualitative risk levels

Negligible risk: less than 2 days of natural background exposure

Minimal risk: more than 2 days and up to 1 month of natural background exposure

Very low risk: more than 1 month and up to 8 months of natural background exposure

Low risk: more than 8 months and up to 6 years of natural background exposure

Moderate risk: more than 6 years of natural background exposure

View table showing summary of radiation doses for a variety of procedures

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2018 Radiological Society of North America, Inc.