

Transgender Breast Cancer Screening

For transgender and gender-nonconforming individuals, breast cancer screening recommendations are based on the sex assigned at birth, risk factors, and the use of hormones.

In average-risk transfeminine (male-to-female) individuals 40 years of age or older with 5 or more years of hormone use, digital breast tomosynthesis (DBT) (<https://www.radiologyinfo.org/en/info/tomosynthesis>) or mammographic screening (<https://www.radiologyinfo.org/en/info/mammo>) may be appropriate. In transfeminine individuals at higher than average risk 25 to 30 years of age or older with 5 or more years of hormone use, DBT or mammographic screening is usually appropriate. In average-risk transfeminine individuals with little to no hormone use (less than 5 years), no test is appropriate. In transfeminine individuals at higher than average risk 25 to 30 years of age or older with little to no hormone use, DBT or mammographic screening may be appropriate.

For transmasculine (female-to-male) individuals with bilateral mastectomies ("top surgery") of any age or risk, no test is appropriate. In average-risk transmasculine individuals who are 40 years of age or older with breast reduction or no chest surgery, DBT or mammographic screening is usually appropriate. In intermediate-risk transmasculine individuals 30 years of age or older with breast reduction or no chest surgery, DBT or mammography screening is usually appropriate, and breast ultrasound (<https://www.radiologyinfo.org/en/info/breastus>) or breast MRI (<https://www.radiologyinfo.org/en/info/breastmr>) with and without intravenous contrast may be appropriate.

In high-risk transmasculine individuals who are 25 to 30 years of age or older with breast reduction or no chest surgery, DBT, mammographic screening, or breast MRI with and without contrast are usually appropriate, and breast ultrasound may be appropriate.

For more information, please see the Breast Cancer Screening (<https://www.radiologyinfo.org/en/info/screening-breast>) page.

— By Ryan Lockhart and Aya Kamaya, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple

distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2026 Radiological Society of North America, Inc.